

Antimicrobial susceptibility results

1/1/2020-12/31/2020

Gram positive cocci	<i>Enterococcus faecalis</i>	<i>Enterococcus faecium</i>	MSSA	MRSA	<i>Staphylococcus epidermidis</i>	<i>Staphylococcus lugdunensis</i>	Group B <i>Streptococcus</i>	<i>Streptococcus pneumoniae</i>
Max strains tested	607	46	704	272	97	73	57	49
% Susceptible								
Ampicillin	99	32	-	0	-	-	-	-
Ciprofloxacin	86*	22*	90	41	73	100	-	-
Clindamycin	-	-	82	78	72	87	36	-
Erythromycin	-	-	67**	14**	34**	89**	25	-
Daptomycin	99	-	99	100	91	100	-	-
Linezolid	90	94	99	100	100	-	-	-
Oxacillin/Nafcillin	-	-	100	0	46	87	-	-
Tetracycline	29	-	94***	92***	81***	98***	-	-
Trimethoprim/sulfa	-	-	98	97	67	987	-	-
Vancomycin	99	90	100	100	98	100	-	-
Penicillin (CSF)	-	-	-	-	-	-	-	69
Penicillin (Non-CSF)	-	-	-	-	-	-	-	95
Ceftriaxone (CSF)	-	-	-	-	-	-	-	86
Ceftriaxone (Non-CSF)	-	-	-	-	-	-	-	87
Cefazolin	-	-	100	0	41	86	-	-
Ceftaroline	-	-	-	100	-	-	-	-
For urinary tract infections only								
Nitrofurantoin	99	19	99	99	100	100	-	-

*Should only be used to treat urinary tract infections caused by *Enterococcus* species.

**Susceptibility to erythromycin can be used to infer susceptibility to azithromycin.

***Susceptibility to tetracycline can be used to infer susceptibility to doxycycline.